

**Gregory Hartman, Ph.D.**

*Department of Applied Mathematics*

*Virginia Military Institute*

***Contributing Authors***

**Troy Siemers, Ph.D.**

*Department of Applied Mathematics*

*Virginia Military Institute*

**Brian Heinold, Ph.D.**

*Department of Mathematics and Computer Science*

*Mount Saint Mary's University*

**Dimplekumar Chalishajar, Ph.D.**

*Department of Applied Mathematics*


*Virginia Military Institute*

***Editor***

**Jennifer Bowen, Ph.D.**

*Department of Mathematics and Computer Science*

*The College of Wooster*



Copyright © 2018 Gregory Hartman  
Licensed to the public under Creative Commons  
Attribution-Noncommercial 4.0 International Public  
License

# Contents

<b>Table of Contents</b>	<b>iii</b>
<b>Preface</b>	<b>vii</b>
<b>1 Limits</b>	<b>1</b>
1.1 An Introduction To Limits . . . . .	1
1.2 Epsilon-Delta Definition of a Limit . . . . .	9
1.3 Finding Limits Analytically . . . . .	18
1.4 One Sided Limits . . . . .	30
1.5 Continuity . . . . .	37
1.6 Limits Involving Infinity . . . . .	46
<b>2 Derivatives</b>	<b>59</b>
2.1 Instantaneous Rates of Change: The Derivative . . . . .	59
2.2 Interpretations of the Derivative . . . . .	75
2.3 Basic Differentiation Rules . . . . .	82
2.4 The Product and Quotient Rules . . . . .	89
2.5 The Chain Rule . . . . .	100
2.6 Implicit Differentiation . . . . .	111
2.7 Derivatives of Inverse Functions . . . . .	122
<b>3 The Graphical Behavior of Functions</b>	<b>129</b>
3.1 Extreme Values . . . . .	129
3.2 The Mean Value Theorem . . . . .	137
3.3 Increasing and Decreasing Functions . . . . .	142
3.4 Concavity and the Second Derivative . . . . .	151
3.5 Curve Sketching . . . . .	159
<b>4 Applications of the Derivative</b>	<b>167</b>
4.1 Newton's Method . . . . .	167
4.2 Related Rates . . . . .	174
4.3 Optimization . . . . .	181
4.4 Differentials . . . . .	188

<b>5</b>	<b>Integration</b>	<b>197</b>
5.1	Antiderivatives and Indefinite Integration . . . . .	197
5.2	The Definite Integral . . . . .	207
5.3	Riemann Sums . . . . .	218
5.4	The Fundamental Theorem of Calculus . . . . .	236
5.5	Numerical Integration . . . . .	248
<b>6</b>	<b>Techniques of Antidifferentiation</b>	<b>263</b>
6.1	Substitution . . . . .	263
6.2	Integration by Parts . . . . .	283
6.3	Trigonometric Integrals . . . . .	294
6.4	Trigonometric Substitution . . . . .	304
6.5	Partial Fraction Decomposition . . . . .	313
6.6	Hyperbolic Functions . . . . .	321
6.7	L'Hôpital's Rule . . . . .	332
6.8	Improper Integration . . . . .	341
<b>7</b>	<b>Applications of Integration</b>	<b>353</b>
7.1	Area Between Curves . . . . .	354
7.2	Volume by Cross-Sectional Area; Disk and Washer Methods . . .	362
7.3	The Shell Method . . . . .	370
7.4	Arc Length and Surface Area . . . . .	378
7.5	Work . . . . .	387
7.6	Fluid Forces . . . . .	397
<b>8</b>	<b>Sequences and Series</b>	<b>405</b>
8.1	Sequences . . . . .	405
8.2	Infinite Series . . . . .	419
8.3	Integral and Comparison Tests . . . . .	434
8.4	Ratio and Root Tests . . . . .	443
8.5	Alternating Series and Absolute Convergence . . . . .	449
8.6	Power Series . . . . .	460
8.7	Taylor Polynomials . . . . .	473
8.8	Taylor Series . . . . .	485
<b>9</b>	<b>Curves in the Plane</b>	<b>497</b>
9.1	Conic Sections . . . . .	497
9.2	Parametric Equations . . . . .	511
9.3	Calculus and Parametric Equations . . . . .	521
9.4	Introduction to Polar Coordinates . . . . .	533
9.5	Calculus and Polar Functions . . . . .	546

<b>10 Vectors</b>	<b>559</b>
10.1 Introduction to Cartesian Coordinates in Space . . . . .	559
10.2 An Introduction to Vectors . . . . .	574
10.3 The Dot Product . . . . .	588
10.4 The Cross Product . . . . .	601
10.5 Lines . . . . .	612
10.6 Planes . . . . .	623
<b>11 Vector Valued Functions</b>	<b>631</b>
11.1 Vector-Valued Functions . . . . .	631
11.2 Calculus and Vector-Valued Functions . . . . .	637
11.3 The Calculus of Motion . . . . .	651
11.4 Unit Tangent and Normal Vectors . . . . .	664
11.5 The Arc Length Parameter and Curvature . . . . .	673
<b>12 Functions of Several Variables</b>	<b>683</b>
12.1 Introduction to Multivariable Functions . . . . .	683
12.2 Limits and Continuity of Multivariable Functions . . . . .	690
12.3 Partial Derivatives . . . . .	700
12.4 Differentiability and the Total Differential . . . . .	712
12.5 The Multivariable Chain Rule . . . . .	721
12.6 Directional Derivatives . . . . .	729
12.7 Tangent Lines, Normal Lines, and Tangent Planes . . . . .	739
12.8 Extreme Values . . . . .	749
<b>13 Multiple Integration</b>	<b>759</b>
13.1 Iterated Integrals and Area . . . . .	759
13.2 Double Integration and Volume . . . . .	769
13.3 Double Integration with Polar Coordinates . . . . .	780
13.4 Center of Mass . . . . .	787
13.5 Surface Area . . . . .	799
13.6 Volume Between Surfaces and Triple Integration . . . . .	806
13.7 Triple Integration with Cylindrical and Spherical Coordinates . . . . .	828
<b>14 Vector Analysis</b>	<b>839</b>
14.1 Introduction to Line Integrals . . . . .	840
14.2 Vector Fields . . . . .	850
14.3 Line Integrals over Vector Fields . . . . .	859
14.4 Flow, Flux, Green's Theorem and the Divergence Theorem . . . . .	870
14.5 Parametrized Surfaces and Surface Area . . . . .	880
14.6 Surface Integrals . . . . .	891
14.7 The Divergence Theorem and Stokes' Theorem . . . . .	900
<b>A Solutions To Selected Problems</b>	<b>A.1</b>
<b>Index</b>	<b>A.37</b>

# 6: TECHNIQUES OF ANTIDIFFERENTIATION

---

The previous chapter introduced the antiderivative and connected it to signed areas under a curve through the Fundamental Theorem of Calculus. The next chapter explores more applications of definite integrals than just area. As evaluating definite integrals will become important, we will want to find antiderivatives of a variety of functions.

This chapter is devoted to exploring techniques of antidifferentiation. While not every function has an antiderivative in terms of elementary functions (a concept introduced in the section on Numerical Integration), we can still find antiderivatives of a wide variety of functions.

## 6.1 Substitution

We motivate this section with an example. Let  $f(x) = (x^2 + 3x - 5)^{10}$ . We can compute  $f'(x)$  using the Chain Rule. It is:

$$f'(x) = 10(x^2 + 3x - 5)^9 \cdot (2x + 3) = (20x + 30)(x^2 + 3x - 5)^9.$$

Now consider this: What is  $\int (20x + 30)(x^2 + 3x - 5)^9 dx$ ? We have the answer in front of us;

$$\int (20x + 30)(x^2 + 3x - 5)^9 dx = (x^2 + 3x - 5)^{10} + C.$$

How would we have evaluated this indefinite integral without starting with  $f(x)$  as we did?

This section explores *integration by substitution*. It allows us to “undo the Chain Rule.” Substitution allows us to evaluate the above integral without knowing the original function first.

The underlying principle is to rewrite a “complicated” integral of the form  $\int f(x) dx$  as a not-so-complicated integral  $\int h(u) du$ . We’ll formally establish later how this is done. First, consider again our introductory indefinite integral,  $\int (20x + 30)(x^2 + 3x - 5)^9 dx$ . Arguably the most “complicated” part of the integrand is  $(x^2 + 3x - 5)^9$ . We wish to make this simpler; we do so through a substitution. Let  $u = x^2 + 3x - 5$ . Thus

$$(x^2 + 3x - 5)^9 = u^9.$$

We have established  $u$  as a function of  $x$ , so now consider the differential of  $u$ :

$$du = (2x + 3)dx.$$

Keep in mind that  $(2x+3)$  and  $dx$  are multiplied; the  $dx$  is not “just sitting there.”

Return to the original integral and do some substitutions through algebra:

$$\begin{aligned} \int (20x + 30)(x^2 + 3x - 5)^9 dx &= \int 10(2x + 3)(x^2 + 3x - 5)^9 dx \\ &= \int 10 \underbrace{(x^2 + 3x - 5)^9}_u \underbrace{(2x + 3) dx}_{du} \\ &= \int 10u^9 du \\ &= u^{10} + C \quad (\text{replace } u \text{ with } x^2 + 3x - 5) \\ &= (x^2 + 3x - 5)^{10} + C \end{aligned}$$

One might well look at this and think “I (sort of) followed how that worked, but I could never come up with that on my own,” but the process is learnable. This section contains numerous examples through which the reader will gain understanding and mathematical maturity enabling them to regard substitution as a natural tool when evaluating integrals.

We stated before that integration by substitution “undoes” the Chain Rule. Specifically, let  $F(x)$  and  $g(x)$  be differentiable functions and consider the derivative of their composition:

$$\frac{d}{dx}(F(g(x))) = F'(g(x))g'(x).$$

Thus

$$\int F'(g(x))g'(x) dx = F(g(x)) + C.$$

Integration by substitution works by recognizing the “inside” function  $g(x)$  and replacing it with a variable. By setting  $u = g(x)$ , we can rewrite the derivative as

$$\frac{d}{dx}(F(u)) = F'(u)u'.$$

Since  $du = g'(x)dx$ , we can rewrite the above integral as

$$\int F'(g(x))g'(x) dx = \int F'(u)du = F(u) + C = F(g(x)) + C.$$

This concept is important so we restate it in the context of a theorem.

Notes:

**Theorem 6.1.1**    **Integration by Substitution**

Let  $F$  and  $g$  be differentiable functions, where the range of  $g$  is an interval  $I$  contained in the domain of  $F$ . Then

$$\int F'(g(x))g'(x) dx = F(g(x)) + C.$$

If  $u = g(x)$ , then  $du = g'(x)dx$  and

$$\int F'(g(x))g'(x) dx = \int F'(u) du = F(u) + C = F(g(x)) + C.$$

The point of substitution is to make the integration step easy. Indeed, the step  $\int F'(u) du = F(u) + C$  looks easy, as the antiderivative of the derivative of  $F$  is just  $F$ , plus a constant. The “work” involved is making the proper substitution. There is not a step-by-step process that one can memorize; rather, experience will be one’s guide. To gain experience, we now embark on many examples.

**Example 6.1.1**    **Integrating by substitution**

Evaluate  $\int x \sin(x^2 + 5) dx$ .

**SOLUTION**    Knowing that substitution is related to the Chain Rule, we choose to let  $u$  be the “inside” function of  $\sin(x^2 + 5)$ . (This is not *always* a good choice, but it is often the best place to start.)

Let  $u = x^2 + 5$ , hence  $du = 2x dx$ . The integrand has an  $x dx$  term, but not a  $2x dx$  term. (Recall that multiplication is commutative, so the  $x$  does not physically have to be next to  $dx$  for there to be an  $x dx$  term.) We can divide both sides of the  $du$  expression by 2:

$$du = 2x dx \quad \Rightarrow \quad \frac{1}{2} du = x dx.$$

We can now substitute.

$$\begin{aligned} \int x \sin(x^2 + 5) dx &= \int \underbrace{\sin(x^2 + 5)}_u \underbrace{x dx}_{\frac{1}{2} du} \\ &= \int \frac{1}{2} \sin u du \end{aligned}$$

---

Notes:



$$\begin{aligned}
 &= -\frac{1}{2} \cos u + C \quad (\text{now replace } u \text{ with } x^2 + 5) \\
 &= -\frac{1}{2} \cos(x^2 + 5) + C.
 \end{aligned}$$

Thus  $\int x \sin(x^2 + 5) dx = -\frac{1}{2} \cos(x^2 + 5) + C$ . We can check our work by evaluating the derivative of the right hand side.

**Example 6.1.2 Integrating by substitution**

Evaluate  $\int \cos(5x) dx$ .

**SOLUTION** Again let  $u$  replace the “inside” function. Letting  $u = 5x$ , we have  $du = 5dx$ . Since our integrand does not have a  $5dx$  term, we can divide the previous equation by 5 to obtain  $\frac{1}{5}du = dx$ . We can now substitute.

$$\begin{aligned}
 \int \cos(5x) dx &= \int \underbrace{\cos(5x)}_u \underbrace{dx}_{\frac{1}{5}du} \\
 &= \int \frac{1}{5} \cos u du \\
 &= \frac{1}{5} \sin u + C \\
 &= \frac{1}{5} \sin(5x) + C.
 \end{aligned}$$

We can again check our work through differentiation.

The previous example exhibited a common, and simple, type of substitution. The “inside” function was a linear function (in this case,  $y = 5x$ ). When the inside function is linear, the resulting integration is very predictable, outlined here.

**Key Idea 6.1.1 Substitution With A Linear Function**

Consider  $\int F'(ax + b) dx$ , where  $a \neq 0$  and  $b$  are constants. Letting  $u = ax + b$  gives  $du = a \cdot dx$ , leading to the result

$$\int F'(ax + b) dx = \frac{1}{a} F(ax + b) + C.$$

Thus  $\int \sin(7x - 4) dx = -\frac{1}{7} \cos(7x - 4) + C$ . Our next example can use Key Idea 6.1.1, but we will only employ it after going through all of the steps.

---

Notes:

**Example 6.1.3** Integrating by substituting a linear function

Evaluate  $\int \frac{7}{-3x+1} dx$ .

**SOLUTION** View the integrand as the composition of functions  $f(g(x))$ , where  $f(x) = 7/x$  and  $g(x) = -3x + 1$ . Employing our understanding of substitution, we let  $u = -3x + 1$ , the inside function. Thus  $du = -3dx$ . The integrand lacks a  $-3$ ; hence divide the previous equation by  $-3$  to obtain  $-du/3 = dx$ . We can now evaluate the integral through substitution.

$$\begin{aligned} \int \frac{7}{-3x+1} dx &= \int \frac{7}{u} \frac{du}{-3} \\ &= \frac{-7}{3} \int \frac{du}{u} \\ &= \frac{-7}{3} \ln |u| + C \\ &= -\frac{7}{3} \ln |-3x+1| + C. \end{aligned}$$

Using Key Idea 6.1.1 is faster, recognizing that  $u$  is linear and  $a = -3$ . One may want to continue writing out all the steps until they are comfortable with this particular shortcut.

Not all integrals that benefit from substitution have a clear “inside” function. Several of the following examples will demonstrate ways in which this occurs.

**Example 6.1.4** Integrating by substitution

Evaluate  $\int \sin x \cos x dx$ .

**SOLUTION** There is not a composition of function here to exploit; rather, just a product of functions. Do not be afraid to experiment; when given an integral to evaluate, it is often beneficial to think “If I let  $u$  be *this*, then  $du$  must be *that* ...” and see if this helps simplify the integral at all.

In this example, let’s set  $u = \sin x$ . Then  $du = \cos x dx$ , which we have as part of the integrand! The substitution becomes very straightforward:

$$\begin{aligned} \int \sin x \cos x dx &= \int u du \\ &= \frac{1}{2} u^2 + C \\ &= \frac{1}{2} \sin^2 x + C. \end{aligned}$$

---

Notes:

One would do well to ask “What would happen if we let  $u = \cos x$ ?” The result is just as easy to find, yet looks very different. The challenge to the reader is to evaluate the integral letting  $u = \cos x$  and discover why the answer is the same, yet looks different.

Our examples so far have required “basic substitution.” The next example demonstrates how substitutions can be made that often strike the new learner as being “nonstandard.”

**Example 6.1.5 Integrating by substitution**

Evaluate  $\int x\sqrt{x+3} \, dx$ .

**SOLUTION** Recognizing the composition of functions, set  $u = x + 3$ . Then  $du = dx$ , giving what seems initially to be a simple substitution. But at this stage, we have:

$$\int x\sqrt{x+3} \, dx = \int x\sqrt{u} \, du.$$

We cannot evaluate an integral that has both an  $x$  and an  $u$  in it. We need to convert the  $x$  to an expression involving just  $u$ .

Since we set  $u = x + 3$ , we can also state that  $u - 3 = x$ . Thus we can replace  $x$  in the integrand with  $u - 3$ . It will also be helpful to rewrite  $\sqrt{u}$  as  $u^{\frac{1}{2}}$ .

$$\begin{aligned} \int x\sqrt{x+3} \, dx &= \int (u-3)u^{\frac{1}{2}} \, du \\ &= \int (u^{\frac{3}{2}} - 3u^{\frac{1}{2}}) \, du \\ &= \frac{2}{5}u^{\frac{5}{2}} - 2u^{\frac{3}{2}} + C \\ &= \frac{2}{5}(x+3)^{\frac{5}{2}} - 2(x+3)^{\frac{3}{2}} + C. \end{aligned}$$

Checking your work is always a good idea. In this particular case, some algebra will be needed to make one’s answer match the integrand in the original problem.

**Example 6.1.6 Integrating by substitution**

Evaluate  $\int \frac{1}{x \ln x} \, dx$ .

**SOLUTION** This is another example where there does not seem to be an obvious composition of functions. The line of thinking used in Example 6.1.5 is useful here: choose something for  $u$  and consider what this implies  $du$  must

---

Notes:

be. If  $u$  can be chosen such that  $du$  also appears in the integrand, then we have chosen well.

Choosing  $u = 1/x$  makes  $du = -1/x^2 dx$ ; that does not seem helpful. However, setting  $u = \ln x$  makes  $du = 1/x dx$ , which is part of the integrand. Thus:

$$\begin{aligned} \int \frac{1}{x \ln x} dx &= \int \underbrace{\frac{1}{\ln x}}_u \underbrace{\frac{1}{x} dx}_{du} \\ &= \int \frac{1}{u} du \\ &= \ln |u| + C \\ &= \ln |\ln x| + C. \end{aligned}$$

The final answer is interesting; the natural log of the natural log. Take the derivative to confirm this answer is indeed correct.

## Integrals Involving Trigonometric Functions

Section 6.3 delves deeper into integrals of a variety of trigonometric functions; here we use substitution to establish a foundation that we will build upon.

The next three examples will help fill in some missing pieces of our antiderivative knowledge. We know the antiderivatives of the sine and cosine functions; what about the other standard functions tangent, cotangent, secant and cosecant? We discover these next.

### Example 6.1.7 Integration by substitution: antiderivatives of $\tan x$

Evaluate  $\int \tan x dx$ .

**SOLUTION** The previous paragraph established that we did not know the antiderivatives of tangent, hence we must assume that we have learned something in this section that can help us evaluate this indefinite integral.

Rewrite  $\tan x$  as  $\sin x / \cos x$ . While the presence of a composition of functions may not be immediately obvious, recognize that  $\cos x$  is “inside” the  $1/x$  function. Therefore, we see if setting  $u = \cos x$  returns usable results. We have

---

Notes:

that  $du = -\sin x \, dx$ , hence  $-du = \sin x \, dx$ . We can integrate:

$$\begin{aligned}\int \tan x \, dx &= \int \frac{\sin x}{\cos x} \, dx \\ &= \int \underbrace{\frac{1}{\cos x}}_u \underbrace{\sin x \, dx}_{-du} \\ &= \int \frac{-1}{u} \, du \\ &= -\ln |u| + C \\ &= -\ln |\cos x| + C.\end{aligned}$$

Some texts prefer to bring the  $-1$  inside the logarithm as a power of  $\cos x$ , as in:

$$\begin{aligned}-\ln |\cos x| + C &= \ln |(\cos x)^{-1}| + C \\ &= \ln \left| \frac{1}{\cos x} \right| + C \\ &= \ln |\sec x| + C.\end{aligned}$$

Thus the result they give is  $\int \tan x \, dx = \ln |\sec x| + C$ . These two answers are equivalent.

**Example 6.1.8 Integrating by substitution: antiderivatives of  $\sec x$**

Evaluate  $\int \sec x \, dx$ .

**SOLUTION** This example employs a wonderful trick: multiply the integrand by “1” so that we see how to integrate more clearly. In this case, we write “1” as

$$1 = \frac{\sec x + \tan x}{\sec x + \tan x}.$$

This may seem like it came out of left field, but it works beautifully. Consider:

$$\begin{aligned}\int \sec x \, dx &= \int \sec x \cdot \frac{\sec x + \tan x}{\sec x + \tan x} \, dx \\ &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} \, dx.\end{aligned}$$

---

Notes:

Now let  $u = \sec x + \tan x$ ; this means  $du = (\sec x \tan x + \sec^2 x) dx$ , which is our numerator. Thus:

$$\begin{aligned} &= \int \frac{du}{u} \\ &= \ln |u| + C \\ &= \ln |\sec x + \tan x| + C. \end{aligned}$$

We can use similar techniques to those used in Examples 6.1.7 and 6.1.8 to find antiderivatives of  $\cot x$  and  $\csc x$  (which the reader can explore in the exercises.) We summarize our results here.

**Theorem 6.1.2 Antiderivatives of Trigonometric Functions**

- |  |   |
|--|---|
| 1. $\int \sin x \, dx = -\cos x + C$       | 4. $\int \csc x \, dx = -\ln  \csc x + \cot x  + C$ |
| 2. $\int \cos x \, dx = \sin x + C$        | 5. $\int \sec x \, dx = \ln  \sec x + \tan x  + C$  |
| 3. $\int \tan x \, dx = -\ln  \cos x  + C$ | 6. $\int \cot x \, dx = \ln  \sin x  + C$           |

We explore one more common trigonometric integral.

**Example 6.1.9 Integration by substitution: powers of  $\cos x$  and  $\sin x$**

Evaluate  $\int \cos^2 x \, dx$ .

**SOLUTION** We have a composition of functions as  $\cos^2 x = (\cos x)^2$ . However, setting  $u = \cos x$  means  $du = -\sin x \, dx$ , which we do not have in the integral. Another technique is needed.

The process we'll employ is to use a Power Reducing formula for  $\cos^2 x$  (perhaps consult the back of this text for this formula), which states

$$\cos^2 x = \frac{1 + \cos(2x)}{2}.$$

The right hand side of this equation is not difficult to integrate. We have:

$$\begin{aligned} \int \cos^2 x \, dx &= \int \frac{1 + \cos(2x)}{2} \, dx \\ &= \int \left( \frac{1}{2} + \frac{1}{2} \cos(2x) \right) \, dx. \end{aligned}$$

Notes:

Now use Key Idea 6.1.1:

$$\begin{aligned}
 &= \frac{1}{2}x + \frac{1}{2} \frac{\sin(2x)}{2} + C \\
 &= \frac{1}{2}x + \frac{\sin(2x)}{4} + C.
 \end{aligned}$$

We'll make significant use of this power-reducing technique in future sections.

### Simplifying the Integrand

It is common to be reluctant to manipulate the integrand of an integral; at first, our grasp of integration is tenuous and one may think that working with the integrand will improperly change the results. Integration by substitution works using a different logic: as long as *equality* is maintained, the integrand can be manipulated so that its *form* is easier to deal with. The next two examples demonstrate common ways in which using algebra first makes the integration easier to perform.

#### Example 6.1.10 Integration by substitution: simplifying first

Evaluate  $\int \frac{x^3 + 4x^2 + 8x + 5}{x^2 + 2x + 1} dx$ .

**SOLUTION** One may try to start by setting  $u$  equal to either the numerator or denominator; in each instance, the result is not workable.

When dealing with rational functions (i.e., quotients made up of polynomial functions), it is an almost universal rule that everything works better when the degree of the numerator is less than the degree of the denominator. Hence we use polynomial division.

We skip the specifics of the steps, but note that when  $x^2 + 2x + 1$  is divided into  $x^3 + 4x^2 + 8x + 5$ , it goes in  $x + 2$  times with a remainder of  $3x + 3$ . Thus

$$\frac{x^3 + 4x^2 + 8x + 5}{x^2 + 2x + 1} = x + 2 + \frac{3x + 3}{x^2 + 2x + 1}.$$

Integrating  $x + 2$  is simple. The fraction can be integrated by setting  $u = x^2 + 2x + 1$ , giving  $du = (2x + 2) dx$ . This is very similar to the numerator. Note that

---

Notes:

$du/2 = (x + 1) dx$  and then consider the following:

$$\begin{aligned} \int \frac{x^3 + 4x^2 + 8x + 5}{x^2 + 2x + 1} dx &= \int \left( x + 2 + \frac{3x + 3}{x^2 + 2x + 1} \right) dx \\ &= \int (x + 2) dx + \int \frac{3(x + 1)}{x^2 + 2x + 1} dx \\ &= \frac{1}{2}x^2 + 2x + C_1 + \int \frac{3}{u} \frac{du}{2} \\ &= \frac{1}{2}x^2 + 2x + C_1 + \frac{3}{2} \ln |u| + C_2 \\ &= \frac{1}{2}x^2 + 2x + \frac{3}{2} \ln |x^2 + 2x + 1| + C. \end{aligned}$$

In some ways, we “lucked out” in that after dividing, substitution was able to be done. In later sections we’ll develop techniques for handling rational functions where substitution is not directly feasible.

**Example 6.1.11 Integration by alternate methods**

Evaluate  $\int \frac{x^2 + 2x + 3}{\sqrt{x}} dx$  with, and without, substitution.

**SOLUTION** We already know how to integrate this particular example. Rewrite  $\sqrt{x}$  as  $x^{1/2}$  and simplify the fraction:

$$\frac{x^2 + 2x + 3}{x^{1/2}} = x^{3/2} + 2x^{1/2} + 3x^{-1/2}.$$

We can now integrate using the Power Rule:

$$\begin{aligned} \int \frac{x^2 + 2x + 3}{x^{1/2}} dx &= \int \left( x^{3/2} + 2x^{1/2} + 3x^{-1/2} \right) dx \\ &= \frac{2}{5}x^{5/2} + \frac{4}{3}x^{3/2} + 6x^{1/2} + C \end{aligned}$$

This is a perfectly fine approach. We demonstrate how this can also be solved using substitution as its implementation is rather clever.

Let  $u = \sqrt{x} = x^{1/2}$ ; therefore

$$du = \frac{1}{2}x^{-1/2}dx = \frac{1}{2\sqrt{x}} dx \quad \Rightarrow \quad 2du = \frac{1}{\sqrt{x}} dx.$$

This gives us  $\int \frac{x^2 + 2x + 3}{\sqrt{x}} dx = \int (x^2 + 2x + 3) \cdot 2 du$ . What are we to do with the other  $x$  terms? Since  $u = x^{1/2}$ ,  $u^2 = x$ , etc. We can then replace  $x^2$  and

Notes:



$x$  with appropriate powers of  $u$ . We thus have

$$\begin{aligned}\int \frac{x^2 + 2x + 3}{\sqrt{x}} dx &= \int (x^2 + 2x + 3) \cdot 2 du \\ &= \int 2(u^4 + 2u^2 + 3) du \\ &= \frac{2}{5}u^5 + \frac{4}{3}u^3 + 6u + C \\ &= \frac{2}{5}x^{\frac{5}{2}} + \frac{4}{3}x^{\frac{3}{2}} + 6x^{\frac{1}{2}} + C,\end{aligned}$$

which is obviously the same answer we obtained before. In this situation, substitution is arguably more work than our other method. The fantastic thing is that it works. It demonstrates how flexible integration is.

### Substitution and Inverse Trigonometric Functions

When studying derivatives of inverse functions, we learned that

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1 + x^2}.$$

Applying the Chain Rule to this is not difficult; for instance,

$$\frac{d}{dx}(\tan^{-1} 5x) = \frac{5}{1 + 25x^2}.$$

We now explore how Substitution can be used to “undo” certain derivatives that are the result of the Chain Rule applied to Inverse Trigonometric functions. We begin with an example.

#### Example 6.1.12 Integrating by substitution: inverse trigonometric functions

Evaluate  $\int \frac{1}{25 + x^2} dx$ .

**SOLUTION** The integrand looks similar to the derivative of the arctangent function. Note:

$$\begin{aligned}\frac{1}{25 + x^2} &= \frac{1}{25(1 + \frac{x^2}{25})} \\ &= \frac{1}{25(1 + (\frac{x}{5})^2)} \\ &= \frac{1}{25} \frac{1}{1 + (\frac{x}{5})^2}.\end{aligned}$$

---

Notes:

Thus

$$\int \frac{1}{25 + x^2} dx = \frac{1}{25} \int \frac{1}{1 + \left(\frac{x}{5}\right)^2} dx.$$

This can be integrated using Substitution. Set  $u = x/5$ , hence  $du = dx/5$  or  $dx = 5du$ . Thus

$$\begin{aligned} \int \frac{1}{25 + x^2} dx &= \frac{1}{25} \int \frac{1}{1 + \left(\frac{x}{5}\right)^2} dx \\ &= \frac{1}{5} \int \frac{1}{1 + u^2} du \\ &= \frac{1}{5} \tan^{-1} u + C \\ &= \frac{1}{5} \tan^{-1} \left(\frac{x}{5}\right) + C \end{aligned}$$

Example 6.1.12 demonstrates a general technique that can be applied to other integrands that result in inverse trigonometric functions. The results are summarized here.

**Theorem 6.1.3 Integrals Involving Inverse Trigonometric Functions**

Let  $a > 0$ .

1.  $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a}\right) + C$
2.  $\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \left(\frac{x}{a}\right) + C$
3.  $\int \frac{1}{x\sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1} \left(\frac{|x|}{a}\right) + C$

Let's practice using Theorem 6.1.3.

**Example 6.1.13 Integrating by substitution: inverse trigonometric functions**

Evaluate the given indefinite integrals.

$$1. \int \frac{1}{9 + x^2} dx, \quad 2. \int \frac{1}{x\sqrt{x^2 - \frac{1}{100}}} dx \quad 3. \int \frac{1}{\sqrt{5 - x^2}} dx.$$

Notes:

**SOLUTION** Each can be answered using a straightforward application of Theorem 6.1.3.

$$1. \int \frac{1}{9+x^2} dx = \frac{1}{3} \tan^{-1} \frac{x}{3} + C, \text{ as } a = 3.$$

$$2. \int \frac{1}{x\sqrt{x^2 - \frac{1}{100}}} dx = 10 \sec^{-1} 10x + C, \text{ as } a = \frac{1}{10}.$$

$$3. \int \frac{1}{\sqrt{5-x^2}} = \sin^{-1} \frac{x}{\sqrt{5}} + C, \text{ as } a = \sqrt{5}.$$

Most applications of Theorem 6.1.3 are not as straightforward. The next examples show some common integrals that can still be approached with this theorem.

**Example 6.1.14 Integrating by substitution: completing the square**

Evaluate  $\int \frac{1}{x^2 - 4x + 13} dx$ .

**SOLUTION** Initially, this integral seems to have nothing in common with the integrals in Theorem 6.1.3. As it lacks a square root, it almost certainly is not related to arcsine or arcsecant. It is, however, related to the arctangent function.

We see this by *completing the square* in the denominator. We give a brief reminder of the process here.

Start with a quadratic with a leading coefficient of 1. It will have the form of  $x^2 + bx + c$ . Take  $1/2$  of  $b$ , square it, and add/subtract it back into the expression. I.e.,

$$\begin{aligned} x^2 + bx + c &= x^2 + bx + \underbrace{\frac{b^2}{4} - \frac{b^2}{4}}_{(x+b/2)^2} + c \\ &= \left(x + \frac{b}{2}\right)^2 + c - \frac{b^2}{4} \end{aligned}$$

In our example, we take half of  $-4$  and square it, getting 4. We add/subtract it into the denominator as follows:

$$\begin{aligned} \frac{1}{x^2 - 4x + 13} &= \frac{1}{\underbrace{x^2 - 4x + 4}_{(x-2)^2} - 4 + 13} \\ &= \frac{1}{(x-2)^2 + 9} \end{aligned}$$

Notes:

We can now integrate this using the arctangent rule. Technically, we need to substitute first with  $u = x - 2$ , but we can employ Key Idea 6.1.1 instead. Thus we have

$$\int \frac{1}{x^2 - 4x + 13} dx = \int \frac{1}{(x-2)^2 + 9} dx = \frac{1}{3} \tan^{-1} \frac{x-2}{3} + C.$$

### Example 6.1.15 Integrals requiring multiple methods

Evaluate  $\int \frac{4-x}{\sqrt{16-x^2}} dx$ .

**SOLUTION** This integral requires two different methods to evaluate it. We get to those methods by splitting up the integral:

$$\int \frac{4-x}{\sqrt{16-x^2}} dx = \int \frac{4}{\sqrt{16-x^2}} dx - \int \frac{x}{\sqrt{16-x^2}} dx.$$

The first integral is handled using a straightforward application of Theorem 6.1.3; the second integral is handled by substitution, with  $u = 16 - x^2$ . We handle each separately.

$$\int \frac{4}{\sqrt{16-x^2}} dx = 4 \sin^{-1} \frac{x}{4} + C.$$

$\int \frac{x}{\sqrt{16-x^2}} dx$ : Set  $u = 16 - x^2$ , so  $du = -2x dx$  and  $x dx = -du/2$ . We have

$$\begin{aligned} \int \frac{x}{\sqrt{16-x^2}} dx &= \int \frac{-du/2}{\sqrt{u}} \\ &= -\frac{1}{2} \int \frac{1}{\sqrt{u}} du \\ &= -\sqrt{u} + C \\ &= -\sqrt{16-x^2} + C. \end{aligned}$$

Combining these together, we have

$$\int \frac{4-x}{\sqrt{16-x^2}} dx = 4 \sin^{-1} \frac{x}{4} + \sqrt{16-x^2} + C.$$

## Substitution and Definite Integration

This section has focused on evaluating indefinite integrals as we are learning a new technique for finding antiderivatives. However, much of the time integration is used in the context of a definite integral. Definite integrals that require substitution can be calculated using the following workflow:

---

Notes:

1. Start with a definite integral  $\int_a^b f(x) dx$  that requires substitution.
2. Ignore the bounds; use substitution to evaluate  $\int f(x) dx$  and find an antiderivative  $F(x)$ .
3. Evaluate  $F(x)$  at the bounds; that is, evaluate  $F(x)\Big|_a^b = F(b) - F(a)$ .

This workflow works fine, but substitution offers an alternative that is powerful and amazing (and a little time saving).

At its heart, (using the notation of Theorem 6.1.1) substitution converts integrals of the form  $\int F'(g(x))g'(x) dx$  into an integral of the form  $\int F'(u) du$  with the substitution of  $u = g(x)$ . The following theorem states how the bounds of a definite integral can be changed as the substitution is performed.

**Theorem 6.1.4 Substitution with Definite Integrals**

Let  $F$  and  $g$  be differentiable functions, where the range of  $g$  is an interval  $I$  that is contained in the domain of  $F$ . Then

$$\int_a^b F'(g(x))g'(x) dx = \int_{g(a)}^{g(b)} F'(u) du.$$

In effect, Theorem 6.1.4 states that once you convert to integrating with respect to  $u$ , you do not need to switch back to evaluating with respect to  $x$ . A few examples will help one understand.

**Example 6.1.16 Definite integrals and substitution: changing the bounds**

Evaluate  $\int_0^2 \cos(3x - 1) dx$  using Theorem 6.1.4.

**SOLUTION** Observing the composition of functions, let  $u = 3x - 1$ , hence  $du = 3dx$ . As  $3dx$  does not appear in the integrand, divide the latter equation by 3 to get  $du/3 = dx$ .

By setting  $u = 3x - 1$ , we are implicitly stating that  $g(x) = 3x - 1$ . Theorem 6.1.4 states that the new lower bound is  $g(0) = -1$ ; the new upper bound is

---

Notes:

$g(2) = 5$ . We now evaluate the definite integral:

$$\begin{aligned}\int_0^2 \cos(3x - 1) dx &= \int_{-1}^5 \cos u \frac{du}{3} \\ &= \frac{1}{3} \sin u \Big|_{-1}^5 \\ &= \frac{1}{3} (\sin 5 - \sin(-1)) \approx -0.039.\end{aligned}$$

Notice how once we converted the integral to be in terms of  $u$ , we never went back to using  $x$ .

The graphs in Figure 6.1.1 tell more of the story. In (a) the area defined by the original integrand is shaded, whereas in (b) the area defined by the new integrand is shaded. In this particular situation, the areas look very similar; the new region is “shorter” but “wider,” giving the same area.

**Example 6.1.17** Definite integrals and substitution: changing the bounds

Evaluate  $\int_0^{\pi/2} \sin x \cos x dx$  using Theorem 6.1.4.

**SOLUTION** We saw the corresponding indefinite integral in Example 6.1.4. In that example we set  $u = \sin x$  but stated that we could have let  $u = \cos x$ . For variety, we do the latter here.


Let  $u = g(x) = \cos x$ , giving  $du = -\sin x dx$  and hence  $\sin x dx = -du$ . The new upper bound is  $g(\pi/2) = 0$ ; the new lower bound is  $g(0) = 1$ . Note how the lower bound is actually larger than the upper bound now. We have

$$\begin{aligned}\int_0^{\pi/2} \sin x \cos x dx &= \int_1^0 -u du \quad (\text{switch bounds \& change sign}) \\ &= \int_0^1 u du \\ &= \frac{1}{2} u^2 \Big|_0^1 = 1/2.\end{aligned}$$


In Figure 6.1.2 we have again graphed the two regions defined by our definite integrals. Unlike the previous example, they bear no resemblance to each other. However, Theorem 6.1.4 guarantees that they have the same area.

Integration by substitution is a powerful and useful integration technique. The next section introduces another technique, called Integration by Parts. As substitution “undoes” the Chain Rule, integration by parts “undoes” the Product Rule. Together, these two techniques provide a strong foundation on which most other integration techniques are based.

Notes:




(a)




(b)

Figure 6.1.1: Graphing the areas defined by the definite integrals of Example 6.1.16.



(a)



(b)

Figure 6.1.2: Graphing the areas defined by the definite integrals of Example 6.1.17.

# Exercises 6.1

## Terms and Concepts

1. Substitution “undoes” what derivative rule?
2. T/F: One can use algebra to rewrite the integrand of an integral to make it easier to evaluate.

## Problems

In Exercises 3 – 14, evaluate the indefinite integral to develop an understanding of Substitution.

3.  $\int 3x^2 (x^3 - 5)^7 dx$
4.  $\int (2x - 5) (x^2 - 5x + 7)^3 dx$
5.  $\int x (x^2 + 1)^8 dx$
6.  $\int (12x + 14) (3x^2 + 7x - 1)^5 dx$
7.  $\int \frac{1}{2x + 7} dx$
8.  $\int \frac{1}{\sqrt{2x + 3}} dx$
9.  $\int \frac{x}{\sqrt{x + 3}} dx$
10.  $\int \frac{x^3 - x}{\sqrt{x}} dx$
11.  $\int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx$
12.  $\int \frac{x^4}{\sqrt{x^5 + 1}} dx$
13.  $\int \frac{\frac{1}{x} + 1}{x^2} dx$
14.  $\int \frac{\ln(x)}{x} dx$

In Exercises 15 – 24, use Substitution to evaluate the indefinite integral involving trigonometric functions.

15.  $\int \sin^2(x) \cos(x) dx$
16.  $\int \cos^3(x) \sin(x) dx$

17.  $\int \cos(3 - 6x) dx$
18.  $\int \sec^2(4 - x) dx$
19.  $\int \sec(2x) dx$
20.  $\int \tan^2(x) \sec^2(x) dx$
21.  $\int x \cos(x^2) dx$
22.  $\int \tan^2(x) dx$
23.  $\int \cot x dx$ . Do not just refer to Theorem 6.1.2 for the answer; justify it through Substitution.
24.  $\int \csc x dx$ . Do not just refer to Theorem 6.1.2 for the answer; justify it through Substitution.

In Exercises 25 – 32, use Substitution to evaluate the indefinite integral involving exponential functions.

25.  $\int e^{3x-1} dx$
26.  $\int e^{x^3} x^2 dx$
27.  $\int e^{x^2-2x+1} (x-1) dx$
28.  $\int \frac{e^x + 1}{e^x} dx$
29.  $\int \frac{e^x}{e^x + 1} dx$
30.  $\int \frac{e^x - e^{-x}}{e^{2x}} dx$
31.  $\int 3^{3x} dx$
32.  $\int 4^{2x} dx$

In Exercises 33 – 36, use Substitution to evaluate the indefinite integral involving logarithmic functions.

33.  $\int \frac{\ln x}{x} dx$
34.  $\int \frac{(\ln x)^2}{x} dx$

35.  $\int \frac{\ln(x^3)}{x} dx$

36.  $\int \frac{1}{x \ln(x^2)} dx$

In Exercises 37 – 42, use Substitution to evaluate the indefinite integral involving rational functions.

37.  $\int \frac{x^2 + 3x + 1}{x} dx$

38.  $\int \frac{x^3 + x^2 + x + 1}{x} dx$

39.  $\int \frac{x^3 - 1}{x + 1} dx$

40.  $\int \frac{x^2 + 2x - 5}{x - 3} dx$

41.  $\int \frac{3x^2 - 5x + 7}{x + 1} dx$

42.  $\int \frac{x^2 + 2x + 1}{x^3 + 3x^2 + 3x} dx$

In Exercises 43 – 52, use Substitution to evaluate the indefinite integral involving inverse trigonometric functions.

43.  $\int \frac{7}{x^2 + 7} dx$

44.  $\int \frac{3}{\sqrt{9 - x^2}} dx$

45.  $\int \frac{14}{\sqrt{5 - x^2}} dx$

46.  $\int \frac{2}{x\sqrt{x^2 - 9}} dx$

47.  $\int \frac{5}{\sqrt{x^4 - 16x^2}} dx$

48.  $\int \frac{x}{\sqrt{1 - x^4}} dx$

49.  $\int \frac{1}{x^2 - 2x + 8} dx$

50.  $\int \frac{2}{\sqrt{-x^2 + 6x + 7}} dx$

51.  $\int \frac{3}{\sqrt{-x^2 + 8x + 9}} dx$

52.  $\int \frac{5}{x^2 + 6x + 34} dx$

In Exercises 53 – 78, evaluate the indefinite integral.

53.  $\int \frac{x^2}{(x^3 + 3)^2} dx$

54.  $\int (3x^2 + 2x)(5x^3 + 5x^2 + 2)^8 dx$

55.  $\int \frac{x}{\sqrt{1 - x^2}} dx$

56.  $\int x^2 \csc^2(x^3 + 1) dx$

57.  $\int \sin(x)\sqrt{\cos(x)} dx$

58.  $\int \sin(5x + 1) dx$

59.  $\int \frac{1}{x - 5} dx$

60.  $\int \frac{7}{3x + 2} dx$

61.  $\int \frac{3x^3 + 4x^2 + 2x - 22}{x^2 + 3x + 5} dx$

62.  $\int \frac{2x + 7}{x^2 + 7x + 3} dx$

63.  $\int \frac{9(2x + 3)}{3x^2 + 9x + 7} dx$

64.  $\int \frac{-x^3 + 14x^2 - 46x - 7}{x^2 - 7x + 1} dx$

65.  $\int \frac{x}{x^4 + 81} dx$

66.  $\int \frac{2}{4x^2 + 1} dx$

67.  $\int \frac{1}{x\sqrt{4x^2 - 1}} dx$

68.  $\int \frac{1}{\sqrt{16 - 9x^2}} dx$

69.  $\int \frac{3x - 2}{x^2 - 2x + 10} dx$

70.  $\int \frac{7 - 2x}{x^2 + 12x + 61} dx$

71.  $\int \frac{x^2 + 5x - 2}{x^2 - 10x + 32} dx$

72.  $\int \frac{x^3}{x^2 + 9} dx$



73.  $\int \frac{x^3 - x}{x^2 + 4x + 9} dx$

74.  $\int \frac{\sin(x)}{\cos^2(x) + 1} dx$

75.  $\int \frac{\cos(x)}{\sin^2(x) + 1} dx$

76.  $\int \frac{\cos(x)}{1 - \sin^2(x)} dx$

77.  $\int \frac{3x - 3}{\sqrt{x^2 - 2x - 6}} dx$

78.  $\int \frac{x - 3}{\sqrt{x^2 - 6x + 8}} dx$

**In Exercises 79 – 86, evaluate the definite integral.**

79.  $\int_1^3 \frac{1}{x - 5} dx$

80.  $\int_2^6 x\sqrt{x - 2} dx$

81.  $\int_{-\pi/2}^{\pi/2} \sin^2 x \cos x dx$

82.  $\int_0^1 2x(1 - x^2)^4 dx$

83.  $\int_{-2}^{-1} (x + 1)e^{x^2 + 2x + 1} dx$

84.  $\int_{-1}^1 \frac{1}{1 + x^2} dx$

85.  $\int_2^4 \frac{1}{x^2 - 6x + 10} dx$

86.  $\int_1^{\sqrt{3}} \frac{1}{\sqrt{4 - x^2}} dx$

## 6.2 Integration by Parts

Here's a simple integral that we can't yet evaluate:

$$\int x \cos x \, dx.$$

It's a simple matter to take the derivative of the integrand using the Product Rule, but there is no Product Rule for integrals. However, this section introduces *Integration by Parts*, a method of integration that is based on the Product Rule for derivatives. It will enable us to evaluate this integral.

The Product Rule says that if  $u$  and  $v$  are functions of  $x$ , then  $(uv)' = u'v + uv'$ . For simplicity, we've written  $u$  for  $u(x)$  and  $v$  for  $v(x)$ . Suppose we integrate both sides with respect to  $x$ . This gives

$$\int (uv)' \, dx = \int (u'v + uv') \, dx.$$

By the Fundamental Theorem of Calculus, the left side integrates to  $uv$ . The right side can be broken up into two integrals, and we have

$$uv = \int u'v \, dx + \int uv' \, dx.$$

Solving for the second integral we have

$$\int uv' \, dx = uv - \int u'v \, dx.$$

Using differential notation, we can write  $du = u'(x)dx$  and  $dv = v'(x)dx$  and the expression above can be written as follows:

$$\int u \, dv = uv - \int v \, du.$$

This is the Integration by Parts formula. For reference purposes, we state this in a theorem.

### Theorem 6.2.1 Integration by Parts

Let  $u$  and  $v$  be differentiable functions of  $x$  on an interval  $I$  containing  $a$  and  $b$ . Then

$$\int u \, dv = uv - \int v \, du,$$

and

$$\int_{x=a}^{x=b} u \, dv = uv \Big|_a^b - \int_{x=a}^{x=b} v \, du.$$

---

Notes:

Let's try an example to understand our new technique.

**Example 6.2.1 Integrating using Integration by Parts**

Evaluate  $\int x \cos x \, dx$ .

**SOLUTION** The key to Integration by Parts is to identify part of the integrand as “ $u$ ” and part as “ $dv$ .” Regular practice will help one make good identifications, and later we will introduce some principles that help. For now, let  $u = x$  and  $dv = \cos x \, dx$ .

It is generally useful to make a small table of these values as done below. Right now we only know  $u$  and  $dv$  as shown on the left of Figure 6.2.1; on the right we fill in the rest of what we need. If  $u = x$ , then  $du = dx$ . Since  $dv = \cos x \, dx$ ,  $v$  is an antiderivative of  $\cos x$ . We choose  $v = \sin x$ .

$$\begin{array}{cc} u = x & v = ? \\ du = ? & dv = \cos x \, dx \end{array} \quad \Rightarrow \quad \begin{array}{cc} u = x & v = \sin x \\ du = dx & dv = \cos x \, dx \end{array}$$

Figure 6.2.1: Setting up Integration by Parts.

Now substitute all of this into the Integration by Parts formula, giving

$$\int x \cos x \, dx = x \sin x - \int \sin x \, dx.$$

We can then integrate  $\sin x$  to get  $-\cos x + C$  and overall our answer is

$$\int x \cos x \, dx = x \sin x + \cos x + C.$$

Note how the antiderivative contains a product,  $x \sin x$ . This product is what makes Integration by Parts necessary.

The example above demonstrates how Integration by Parts works in general. We try to identify  $u$  and  $dv$  in the integral we are given, and the key is that we usually want to choose  $u$  and  $dv$  so that  $du$  is simpler than  $u$  and  $v$  is hopefully not too much more complicated than  $dv$ . This will mean that the integral on the right side of the Integration by Parts formula,  $\int v \, du$  will be simpler to integrate than the original integral  $\int u \, dv$ .

In the example above, we chose  $u = x$  and  $dv = \cos x \, dx$ . Then  $du = dx$  was simpler than  $u$  and  $v = \sin x$  is no more complicated than  $dv$ . Therefore, instead of integrating  $x \cos x \, dx$ , we could integrate  $\sin x \, dx$ , which we knew how to do.

A useful mnemonic for helping to determine  $u$  is “LIATE,” where

L = Logarithmic, I = Inverse Trig., A = Algebraic (polynomials),  
T = Trigonometric, and E = Exponential.

---

Notes:

If the integrand contains both a logarithmic and an algebraic term, in general letting  $u$  be the logarithmic term works best, as indicated by L coming before A in LIATE.

We now consider another example.

**Example 6.2.2 Integrating using Integration by Parts**

Evaluate  $\int xe^x dx$ .

**SOLUTION** The integrand contains an **A**lgebraic term ( $x$ ) and an **E**xponential term ( $e^x$ ). Our mnemonic suggests letting  $u$  be the algebraic term, so we choose  $u = x$  and  $dv = e^x dx$ . Then  $du = dx$  and  $v = e^x$  as indicated by the tables below.

$$\begin{array}{ll} u = x & v = ? \\ du = ? & dv = e^x dx \end{array} \Rightarrow \begin{array}{ll} u = x & v = e^x \\ du = dx & dv = e^x dx \end{array}$$

Figure 6.2.2: Setting up Integration by Parts.

We see  $du$  is simpler than  $u$ , while there is no change in going from  $dv$  to  $v$ . This is good. The Integration by Parts formula gives

$$\int xe^x dx = xe^x - \int e^x dx.$$

The integral on the right is simple; our final answer is

$$\int xe^x dx = xe^x - e^x + C.$$

Note again how the antiderivatives contain a product term.

**Example 6.2.3 Integrating using Integration by Parts**

Evaluate  $\int x^2 \cos x dx$ .

**SOLUTION** The mnemonic suggests letting  $u = x^2$  instead of the trigonometric function, hence  $dv = \cos x dx$ . Then  $du = 2x dx$  and  $v = \sin x$  as shown below.

$$\begin{array}{ll} u = x^2 & v = ? \\ du = ? & dv = \cos x dx \end{array} \Rightarrow \begin{array}{ll} u = x^2 & v = \sin x \\ du = 2x dx & dv = \cos x dx \end{array}$$

Figure 6.2.3: Setting up Integration by Parts.

---

Notes:

The Integration by Parts formula gives

$$\int x^2 \cos x \, dx = x^2 \sin x - \int 2x \sin x \, dx.$$

At this point, the integral on the right is indeed simpler than the one we started with, but to evaluate it, we need to do Integration by Parts again. Here we choose  $u = 2x$  and  $dv = \sin x$  and fill in the rest below.

$$\begin{array}{ll} u = 2x & v = ? \\ du = ? & dv = \sin x \, dx \end{array} \quad \Rightarrow \quad \begin{array}{ll} u = 2x & v = -\cos x \\ du = 2 \, dx & dv = \sin x \, dx \end{array}$$

Figure 6.2.4: Setting up Integration by Parts (again).

$$\int x^2 \cos x \, dx = x^2 \sin x - \left( -2x \cos x - \int -2 \cos x \, dx \right).$$

The integral all the way on the right is now something we can evaluate. It evaluates to  $-2 \sin x$ . Then going through and simplifying, being careful to keep all the signs straight, our answer is

$$\int x^2 \cos x \, dx = x^2 \sin x + 2x \cos x - 2 \sin x + C.$$

#### Example 6.2.4 Integrating using Integration by Parts

Evaluate  $\int e^x \cos x \, dx$ .

**SOLUTION** This is a classic problem. Our mnemonic suggests letting  $u$  be the trigonometric function instead of the exponential. In this particular example, one can let  $u$  be either  $\cos x$  or  $e^x$ ; to demonstrate that we do not have to follow LIATE, we choose  $u = e^x$  and hence  $dv = \cos x \, dx$ . Then  $du = e^x \, dx$  and  $v = \sin x$  as shown below.

$$\begin{array}{ll} u = e^x & v = ? \\ du = ? & dv = \cos x \, dx \end{array} \quad \Rightarrow \quad \begin{array}{ll} u = e^x & v = \sin x \\ du = e^x \, dx & dv = \cos x \, dx \end{array}$$

Figure 6.2.5: Setting up Integration by Parts.

Notice that  $du$  is no simpler than  $u$ , going against our general rule (but bear with us). The Integration by Parts formula yields

$$\int e^x \cos x \, dx = e^x \sin x - \int e^x \sin x \, dx.$$

Notes:

The integral on the right is not much different than the one we started with, so it seems like we have gotten nowhere. Let's keep working and apply Integration by Parts to the new integral, using  $u = e^x$  and  $dv = \sin x \, dx$ . This leads us to the following:

$$\begin{array}{ll} u = e^x & v = ? \\ du = ? & dv = \sin x \, dx \end{array} \quad \Rightarrow \quad \begin{array}{ll} u = e^x & v = -\cos x \\ du = e^x \, dx & dv = \sin x \, dx \end{array}$$

Figure 6.2.6: Setting up Integration by Parts (again).

The Integration by Parts formula then gives:

$$\begin{aligned} \int e^x \cos x \, dx &= e^x \sin x - \left( -e^x \cos x - \int -e^x \cos x \, dx \right) \\ &= e^x \sin x + e^x \cos x - \int e^x \cos x \, dx. \end{aligned}$$

It seems we are back right where we started, as the right hand side contains  $\int e^x \cos x \, dx$ . But this is actually a good thing.

Add  $\int e^x \cos x \, dx$  to both sides. This gives

$$2 \int e^x \cos x \, dx = e^x \sin x + e^x \cos x$$

Now divide both sides by 2:

$$\int e^x \cos x \, dx = \frac{1}{2} (e^x \sin x + e^x \cos x).$$

Simplifying a little and adding the constant of integration, our answer is thus

$$\int e^x \cos x \, dx = \frac{1}{2} e^x (\sin x + \cos x) + C.$$

**Example 6.2.5** Integrating using Integration by Parts: antiderivative of  $\ln x$

Evaluate  $\int \ln x \, dx$ .

**SOLUTION** One may have noticed that we have rules for integrating the familiar trigonometric functions and  $e^x$ , but we have not yet given a rule for integrating  $\ln x$ . That is because  $\ln x$  can't easily be integrated with any of the rules we have learned up to this point. But we can find its antiderivative by a

Notes:

clever application of Integration by Parts. Set  $u = \ln x$  and  $dv = dx$ . This is a good, sneaky trick to learn as it can help in other situations. This determines  $du = (1/x) dx$  and  $v = x$  as shown below.

$$\begin{array}{ll} u = \ln x & v = ? \\ du = ? & dv = dx \end{array} \Rightarrow \begin{array}{ll} u = \ln x & v = x \\ du = 1/x dx & dv = dx \end{array}$$

Figure 6.2.7: Setting up Integration by Parts.

Putting this all together in the Integration by Parts formula, things work out very nicely:

$$\int \ln x dx = x \ln x - \int x \frac{1}{x} dx.$$

The new integral simplifies to  $\int 1 dx$ , which is about as simple as things get. Its integral is  $x + C$  and our answer is

$$\int \ln x dx = x \ln x - x + C.$$

**Example 6.2.6 Integrating using Int. by Parts: antiderivative of  $\arctan x$**

Evaluate  $\int \arctan x dx$ .

**SOLUTION** The same sneaky trick we used above works here. Let  $u = \arctan x$  and  $dv = dx$ . Then  $du = 1/(1+x^2) dx$  and  $v = x$ . The Integration by Parts formula gives

$$\int \arctan x dx = x \arctan x - \int \frac{x}{1+x^2} dx.$$

The integral on the right can be solved by substitution. Taking  $u = 1+x^2$ , we get  $du = 2x dx$ . The integral then becomes

$$\int \arctan x dx = x \arctan x - \frac{1}{2} \int \frac{1}{u} du.$$

The integral on the right evaluates to  $\ln |u| + C$ , which becomes  $\ln(1+x^2) + C$  (we can drop the absolute values as  $1+x^2$  is always positive). Therefore, the answer is

$$\int \arctan x dx = x \arctan x - \frac{1}{2} \ln(1+x^2) + C.$$

---

Notes:

### Substitution Before Integration

When taking derivatives, it was common to employ multiple rules (such as using both the Quotient and the Chain Rules). It should then come as no surprise that some integrals are best evaluated by combining integration techniques. In particular, here we illustrate making an “unusual” substitution first before using Integration by Parts.

#### Example 6.2.7 Integration by Parts after substitution

Evaluate  $\int \cos(\ln x) dx$ .

**SOLUTION** The integrand contains a composition of functions, leading us to think Substitution would be beneficial. Letting  $u = \ln x$ , we have  $du = 1/x dx$ . This seems problematic, as we do not have a  $1/x$  in the integrand. But consider:

$$du = \frac{1}{x} dx \Rightarrow x \cdot du = dx.$$

Since  $u = \ln x$ , we can use inverse functions and conclude that  $x = e^u$ . Therefore we have that

$$\begin{aligned} dx &= x \cdot du \\ &= e^u du. \end{aligned}$$

We can thus replace  $\ln x$  with  $u$  and  $dx$  with  $e^u du$ . Thus we rewrite our integral as

$$\int \cos(\ln x) dx = \int e^u \cos u du.$$

We evaluated this integral in Example 6.2.4. Using the result there, we have:

$$\begin{aligned} \int \cos(\ln x) dx &= \int e^u \cos u du \\ &= \frac{1}{2} e^u (\sin u + \cos u) + C \\ &= \frac{1}{2} e^{\ln x} (\sin(\ln x) + \cos(\ln x)) + C \\ &= \frac{1}{2} x (\sin(\ln x) + \cos(\ln x)) + C. \end{aligned}$$

### Definite Integrals and Integration By Parts

So far we have focused only on evaluating indefinite integrals. Of course, we can use Integration by Parts to evaluate definite integrals as well, as Theorem

Notes:



6.2.1 states. We do so in the next example.

**Example 6.2.8** Definite integration using Integration by Parts

Evaluate  $\int_1^2 x^2 \ln x \, dx$ .

**SOLUTION** Our mnemonic suggests letting  $u = \ln x$ , hence  $dv = x^2 \, dx$ . We then get  $du = (1/x) \, dx$  and  $v = x^3/3$  as shown below.

$$\begin{array}{llll} u = \ln x & v = ? & \Rightarrow & u = \ln x \quad v = x^3/3 \\ du = ? & dv = x^2 \, dx & & du = 1/x \, dx \quad dv = x^2 \, dx \end{array}$$

Figure 6.2.8: Setting up Integration by Parts.

The Integration by Parts formula then gives

$$\begin{aligned} \int_1^2 x^2 \ln x \, dx &= \left. \frac{x^3}{3} \ln x \right|_1^2 - \int_1^2 \frac{x^3}{3} \frac{1}{x} \, dx \\ &= \left. \frac{x^3}{3} \ln x \right|_1^2 - \int_1^2 \frac{x^2}{3} \, dx \\ &= \left. \frac{x^3}{3} \ln x - \frac{x^3}{9} \right|_1^2 \\ &= \left( \frac{x^3}{3} \ln x - \frac{x^3}{9} \right) \Big|_1^2 \\ &= \left( \frac{8}{3} \ln 2 - \frac{8}{9} \right) - \left( \frac{1}{3} \ln 1 - \frac{1}{9} \right) \\ &= \frac{8}{3} \ln 2 - \frac{7}{9} \\ &\approx 1.07. \end{aligned}$$

In general, Integration by Parts is useful for integrating certain products of functions, like  $\int xe^x \, dx$  or  $\int x^3 \sin x \, dx$ . It is also useful for integrals involving logarithms and inverse trigonometric functions.

As stated before, integration is generally more difficult than derivation. We are developing tools for handling a large array of integrals, and experience will tell us when one tool is preferable/necessary over another. For instance, consider the three similar-looking integrals

$$\int xe^x \, dx, \quad \int xe^{x^2} \, dx \quad \text{and} \quad \int xe^{x^3} \, dx.$$

---

Notes:

While the first is calculated easily with Integration by Parts, the second is best approached with Substitution. Taking things one step further, the third integral has no answer in terms of elementary functions, so none of the methods we learn in calculus will get us the exact answer.

Integration by Parts is a very useful method, second only to Substitution. In the following sections of this chapter, we continue to learn other integration techniques. The next section focuses on handling integrals containing trigonometric functions.

---

Notes:

## Exercises 6.2

---

### Terms and Concepts

1. T/F: Integration by Parts is useful in evaluating integrands that contain products of functions.
2. T/F: Integration by Parts can be thought of as the “opposite of the Chain Rule.”
3. For what is “LIATE” useful?
4. T/F: If the integral that results from Integration by Parts appears to also need Integration by Parts, then a mistake was made in the original choice of “ $u$ ”.

### Problems

In Exercises 5 – 34, evaluate the given indefinite integral.

5.  $\int x \sin x \, dx$
6.  $\int x e^{-x} \, dx$
7.  $\int x^2 \sin x \, dx$
8.  $\int x^3 \sin x \, dx$
9.  $\int x e^{x^2} \, dx$
10.  $\int x^3 e^x \, dx$
11.  $\int x e^{-2x} \, dx$
12.  $\int e^x \sin x \, dx$
13.  $\int e^{2x} \cos x \, dx$
14.  $\int e^{2x} \sin(3x) \, dx$
15.  $\int e^{5x} \cos(5x) \, dx$
16.  $\int \sin x \cos x \, dx$
17.  $\int \sin^{-1} x \, dx$
18.  $\int \tan^{-1}(2x) \, dx$
19.  $\int x \tan^{-1} x \, dx$
20.  $\int \sin^{-1} x \, dx$
21.  $\int x \ln x \, dx$
22.  $\int (x - 2) \ln x \, dx$
23.  $\int x \ln(x - 1) \, dx$
24.  $\int x \ln(x^2) \, dx$
25.  $\int x^2 \ln x \, dx$
26.  $\int (\ln x)^2 \, dx$
27.  $\int (\ln(x + 1))^2 \, dx$
28.  $\int x \sec^2 x \, dx$
29.  $\int x \csc^2 x \, dx$
30.  $\int x \sqrt{x - 2} \, dx$
31.  $\int x \sqrt{x^2 - 2} \, dx$
32.  $\int \sec x \tan x \, dx$
33.  $\int x \sec x \tan x \, dx$
34.  $\int x \csc x \cot x \, dx$

In Exercises 35 – 40, evaluate the indefinite integral after first making a substitution.

35.  $\int \sin(\ln x) \, dx$
36.  $\int e^{2x} \cos(e^x) \, dx$

$$37. \int \sin(\sqrt{x}) dx$$

$$38. \int \ln(\sqrt{x}) dx$$

$$39. \int e^{\sqrt{x}} dx$$

$$40. \int e^{\ln x} dx$$

**In Exercises 41 – 49, evaluate the definite integral. Note: the corresponding indefinite integrals appear in Exercises 5 – 13.**

$$41. \int_0^{\pi} x \sin x dx$$

$$42. \int_{-1}^1 xe^{-x} dx$$

$$43. \int_{-\pi/4}^{\pi/4} x^2 \sin x dx$$

$$44. \int_{-\pi/2}^{\pi/2} x^3 \sin x dx$$

$$45. \int_0^{\sqrt{\ln 2}} xe^{x^2} dx$$

$$46. \int_0^1 x^3 e^x dx$$

$$47. \int_1^2 xe^{-2x} dx$$

$$48. \int_0^{\pi} e^x \sin x dx$$

$$49. \int_{-\pi/2}^{\pi/2} e^{2x} \cos x dx$$

55.  $F'(x) = (3x^2 + 1) \frac{1}{x^3 + x}$   
 57.  $F'(x) = 2x(x^2 + 2) - (x + 2)$

### Section 5.5

1. F  
 3. They are superseded by the Trapezoidal Rule; it takes an equal amount of work and is generally more accurate.  
 5. (a)  $3/4$   
 (b)  $2/3$   
 (c)  $2/3$   
 7. (a)  $\frac{1}{4}(1 + \sqrt{2})\pi \approx 1.896$   
 (b)  $\frac{1}{6}(1 + 2\sqrt{2})\pi \approx 2.005$   
 (c) 2  
 9. (a) 38.5781  
 (b)  $147/4 \approx 36.75$   
 (c)  $147/4 \approx 36.75$   
 11. (a) 0  
 (b) 0  
 (c) 0  
 13. Trapezoidal Rule: 0.9006  
 Simpson's Rule: 0.90452  
 15. Trapezoidal Rule: 13.9604  
 Simpson's Rule: 13.9066  
 17. Trapezoidal Rule: 1.1703  
 Simpson's Rule: 1.1873  
 19. Trapezoidal Rule: 1.0803  
 Simpson's Rule: 1.077  
 21. (a)  $n = 161$  (using  $\max(f''(x)) = 1$ )  
 (b)  $n = 12$  (using  $\max(f^{(4)}(x)) = 1$ )  
 23. (a)  $n = 1004$  (using  $\max(f''(x)) = 39$ )  
 (b)  $n = 62$  (using  $\max(f^{(4)}(x)) = 800$ )  
 25. (a) Area is  $30.8667 \text{ cm}^2$ .  
 (b) Area is  $308,667 \text{ yd}^2$ .

## Chapter 6

### Section 6.1

1. Chain Rule.  
 3.  $\frac{1}{8}(x^3 - 5)^8 + C$   
 5.  $\frac{1}{18}(x^2 + 1)^9 + C$   
 7.  $\frac{1}{2} \ln |2x + 7| + C$   
 9.  $\frac{2}{3}(x + 3)^{3/2} - 6(x + 3)^{1/2} + C = \frac{2}{3}(x - 6)\sqrt{x + 3} + C$   
 11.  $2e^{\sqrt{x}} + C$   
 13.  $-\frac{1}{2x^2} - \frac{1}{x} + C$   
 15.  $\frac{\sin^3(x)}{3} + C$   
 17.  $-\frac{1}{6} \sin(3 - 6x) + C$   
 19.  $\frac{1}{2} \ln |\sec(2x) + \tan(2x)| + C$   
 21.  $\frac{\sin(x^2)}{2} + C$

23. The key is to rewrite  $\cot x$  as  $\cos x / \sin x$ , and let  $u = \sin x$ .

25.  $\frac{1}{3}e^{3x-1} + C$   
 27.  $\frac{1}{2}e^{(x-1)^2} + C$   
 29.  $\ln(e^x + 1) + C$   
 31.  $\frac{27^x}{\ln 27} + C$   
 33.  $\frac{1}{2} \ln^2(x) + C$   
 35.  $\frac{3}{2}(\ln x)^2 + C$   
 37.  $\frac{x^2}{2} + 3x + \ln|x| + C$   
 39.  $\frac{x^3}{3} - \frac{x^2}{2} + x - 2 \ln|x + 1| + C$   
 41.  $\frac{3}{2}x^2 - 8x + 15 \ln|x + 1| + C$   
 43.  $\sqrt{7} \tan^{-1}\left(\frac{x}{\sqrt{7}}\right) + C$   
 45.  $14 \sin^{-1}\left(\frac{x}{\sqrt{5}}\right) + C$   
 47.  $\frac{5}{4} \sec^{-1}(|x|/4) + C$   
 49.  $\frac{\tan^{-1}\left(\frac{x-1}{\sqrt{7}}\right)}{\sqrt{7}} + C$   
 51.  $3 \sin^{-1}\left(\frac{x-4}{5}\right) + C$   
 53.  $-\frac{1}{3(x^3+3)} + C$   
 55.  $-\sqrt{1-x^2} + C$   
 57.  $-\frac{2}{3} \cos^{\frac{3}{2}}(x) + C$   
 59.  $\ln|x - 5| + C$   
 61.  $\frac{3x^2}{2} + \ln|x^2 + 3x + 5| - 5x + C$   
 63.  $3 \ln|3x^2 + 9x + 7| + C$   
 65.  $\frac{1}{18} \tan^{-1}\left(\frac{x^2}{9}\right) + C$   
 67.  $\sec^{-1}(|2x|) + C$   
 69.  $\frac{3}{2} \ln|x^2 - 2x + 10| + \frac{1}{3} \tan^{-1}\left(\frac{x-1}{3}\right) + C$   
 71.  $\frac{15}{2} \ln|x^2 - 10x + 32| + x + \frac{41 \tan^{-1}\left(\frac{x-5}{\sqrt{7}}\right)}{\sqrt{7}} + C$   
 73.  $\frac{x^2}{2} + 3 \ln|x^2 + 4x + 9| - 4x + \frac{24 \tan^{-1}\left(\frac{x+2}{\sqrt{5}}\right)}{\sqrt{5}} + C$   
 75.  $\tan^{-1}(\sin(x)) + C$   
 77.  $3\sqrt{x^2 - 2x - 6} + C$   
 79.  $-\ln 2$   
 81.  $2/3$   
 83.  $(1 - e)/2$   
 85.  $\pi/2$

### Section 6.2

1. T  
 3. Determining which functions in the integrand to set equal to "u" and which to set equal to "dv".  
 5.  $\sin x - x \cos x + C$   
 7.  $-x^2 \cos x + 2x \sin x + 2 \cos x + C$   
 9.  $1/2e^{x^2} + C$   
 11.  $-\frac{1}{2}xe^{-2x} - \frac{e^{-2x}}{4} + C$   
 13.  $1/5e^{2x}(\sin x + 2 \cos x) + C$   
 15.  $1/10e^{5x}(\sin(5x) + \cos(5x)) + C$

17.  $\sqrt{1-x^2} + x \sin^{-1}(x) + C$
19.  $\frac{1}{2}x^2 \tan^{-1}(x) - \frac{x}{2} + \frac{1}{2} \tan^{-1}(x) + C$
21.  $\frac{1}{2}x^2 \ln|x| - \frac{x^2}{4} + C$
23.  $-\frac{x^2}{4} + \frac{1}{2}x^2 \ln|x-1| - \frac{x}{2} - \frac{1}{2} \ln|x-1| + C$
25.  $\frac{1}{3}x^3 \ln|x| - \frac{x^3}{9} + C$
27.  $2(x+1) + (x+1)(\ln(x+1))^2 - 2(x+1)\ln(x+1) + C$
29.  $\ln|\sin(x)| - x \cot(x) + C$
31.  $\frac{1}{3}(x^2-2)^{3/2} + C$
33.  $x \sec x - \ln|\sec x + \tan x| + C$
35.  $1/2x(\sin(\ln x) - \cos(\ln x)) + C$
37.  $2 \sin(\sqrt{x}) - 2\sqrt{x} \cos(\sqrt{x}) + C$
39.  $2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}} + C$
41.  $\pi$
43.  $0$
45.  $1/2$
47.  $\frac{3}{4e^2} - \frac{5}{4e^4}$
49.  $1/5(e^\pi + e^{-\pi})$

### Section 6.3

1. F
3. F
5.  $-\frac{1}{5} \cos^5(x) + C$
7.  $\frac{1}{5} \cos^5 x - \frac{1}{3} \cos^3 x + C$
9.  $\frac{1}{11} \sin^{11} x - \frac{2}{9} \sin^9 x + \frac{1}{7} \sin^7 x + C$
11.  $\frac{x}{8} - \frac{1}{32} \sin(4x) + C$
13.  $\frac{1}{2}(-\frac{1}{8} \cos(8x) - \frac{1}{2} \cos(2x)) + C$
15.  $\frac{1}{2}(\frac{1}{4} \sin(4x) - \frac{1}{10} \sin(10x)) + C$
17.  $\frac{1}{2}(\sin(x) + \frac{1}{3} \sin(3x)) + C$
19.  $\frac{\tan^5(x)}{5} + C$
21.  $\frac{\tan^6(x)}{6} + \frac{\tan^4(x)}{4} + C$
23.  $\frac{\sec^5(x)}{5} - \frac{\sec^3(x)}{3} + C$
25.  $\frac{1}{3} \tan^3 x - \tan x + x + C$
27.  $\frac{1}{2}(\sec x \tan x - \ln|\sec x + \tan x|) + C$
29.  $\frac{2}{5}$
31.  $32/315$
33.  $2/3$
35.  $16/15$

### Section 6.4

1. backwards
3. (a)  $\tan^2 \theta + 1 = \sec^2 \theta$   
(b)  $9 \sec^2 \theta$ .
5.  $\frac{1}{2}(x\sqrt{x^2+1} + \ln|\sqrt{x^2+1}+x|) + C$
7.  $\frac{1}{2}(\sin^{-1}x + x\sqrt{1-x^2}) + C$
9.  $\frac{1}{2}x\sqrt{x^2-1} - \frac{1}{2} \ln|x + \sqrt{x^2-1}| + C$

11.  $x\sqrt{x^2+1/4} + \frac{1}{4} \ln|2\sqrt{x^2+1/4}+2x| + C = \frac{1}{2}x\sqrt{4x^2+1} + \frac{1}{4} \ln|\sqrt{4x^2+1}+2x| + C$
13.  $4\left(\frac{1}{2}x\sqrt{x^2-1/16} - \frac{1}{32} \ln|4x + 4\sqrt{x^2-1/16}|\right) + C = \frac{1}{2}x\sqrt{16x^2-1} - \frac{1}{8} \ln|4x + \sqrt{16x^2-1}| + C$
15.  $3 \sin^{-1}\left(\frac{x}{\sqrt{7}}\right) + C$  (Trig. Subst. is not needed)
17.  $\sqrt{x^2-11} - \sqrt{11} \sec^{-1}(x/\sqrt{11}) + C$
19.  $\sqrt{x^2-3} + C$  (Trig. Subst. is not needed)
21.  $-\frac{1}{\sqrt{x^2+9}} + C$  (Trig. Subst. is not needed)
23.  $\frac{1}{18} \frac{x+2}{x^2+4x+13} + \frac{1}{54} \tan^{-1}\left(\frac{x+2}{2}\right) + C$
25.  $\frac{1}{7}\left(-\frac{\sqrt{5-x^2}}{x} - \sin^{-1}(x/\sqrt{5})\right) + C$
27.  $\pi/2$
29.  $2\sqrt{2} + 2 \ln(1 + \sqrt{2})$
31.  $9 \sin^{-1}(1/3) + \sqrt{8}$  Note: the new lower bound is  $\theta = \sin^{-1}(-1/3)$  and the new upper bound is  $\theta = \sin^{-1}(1/3)$ . The final answer comes with recognizing that  $\sin^{-1}(-1/3) = -\sin^{-1}(1/3)$  and that  $\cos(\sin^{-1}(1/3)) = \cos(\sin^{-1}(-1/3)) = \sqrt{8}/3$ .

### Section 6.5

1. rational
3.  $\frac{A}{x} + \frac{B}{x-3}$
5.  $\frac{A}{x-\sqrt{7}} + \frac{B}{x+\sqrt{7}}$
7.  $3 \ln|x-2| + 4 \ln|x+5| + C$
9.  $\frac{1}{3}(\ln|x+2| - \ln|x-2|) + C$
11.  $\ln|x+5| - \frac{2}{x+5} + C$
13.  $\frac{5}{x+1} + 7 \ln|x| + 2 \ln|x+1| + C$
15.  $-\frac{1}{5} \ln|5x-1| + \frac{2}{3} \ln|3x-1| + \frac{3}{7} \ln|7x+3| + C$
17.  $\frac{x^2}{2} + x + \frac{125}{9} \ln|x-5| + \frac{64}{9} \ln|x+4| - \frac{35}{2} + C$
19.  $\frac{1}{6}(-\ln|x^2+2x+3| + 2 \ln|x| - \sqrt{2} \tan^{-1}\left(\frac{x+1}{\sqrt{2}}\right)) + C$
21.  $\ln|3x^2+5x-1| + 2 \ln|x+1| + C$
23.  $\frac{9}{10} \ln|x^2+9| + \frac{1}{5} \ln|x+1| - \frac{4}{15} \tan^{-1}\left(\frac{x}{3}\right) + C$
25.  $3(\ln|x^2-2x+11| + \ln|x-9|) + 3\sqrt{\frac{2}{5}} \tan^{-1}\left(\frac{x-1}{\sqrt{10}}\right) + C$
27.  $\ln(2000/243) \approx 2.108$
29.  $-\pi/4 + \tan^{-1} 3 - \ln(11/9) \approx 0.263$

### Section 6.6

1. Because cosh  $x$  is always positive.
3. 
$$\begin{aligned} \coth^2 x - \operatorname{csch}^2 x &= \left(\frac{e^x + e^{-x}}{e^x - e^{-x}}\right)^2 - \left(\frac{2}{e^x - e^{-x}}\right)^2 \\ &= \frac{(e^{2x} + 2 + e^{-2x}) - (4)}{e^{2x} - 2 + e^{-2x}} \\ &= \frac{e^{2x} - 2 + e^{-2x}}{e^{2x} - 2 + e^{-2x}} \\ &= 1 \end{aligned}$$