

Expressions (and reductions)

$$\begin{array}{ccc}
 2 \cdot 3 + 4 \cdot 5 & \longrightarrow & 2 \cdot 3 + 20 \\
 \downarrow & & \downarrow \\
 6 + 4 \cdot 5 & \longrightarrow & 6 + 20 \longrightarrow 26
 \end{array}$$

$$\underbrace{\underbrace{2 \cdot 3}_6 + \underbrace{4 \cdot 5}_{20}}_{26}$$

$$\begin{aligned}
 2 \cdot 3 + 4 \cdot 5 &= 2 \cdot 3 + 20 \\
 &= 6 + 20 \\
 &= 26
 \end{aligned}$$

$$\begin{aligned}
 2 \cdot 3 + 4 \cdot 5 &= 6 + 4 \cdot 5 \\
 &= 6 + 20 \\
 &= 26
 \end{aligned}$$

Subexpressions:

$$\underbrace{\underbrace{2 \cdot 3}_6 + \underbrace{4 \cdot 5}_{20}}_{26}$$

Expressions with variables:

If $a = 5$ and $b = 2$, then:

$$\underbrace{\underbrace{\underbrace{a}_{5} + \underbrace{b}_{2}}_7 \cdot \underbrace{\underbrace{a}_{5} - \underbrace{b}_{2}}_3}_{21}$$

If $a = 10$ and $b = 1$, then:

$$\underbrace{\underbrace{\underbrace{a}_{10} + \underbrace{b}_{1}}_{11} \cdot \underbrace{\underbrace{a}_{10} - \underbrace{b}_{1}}_9}_{99}$$

We know – by algebra, which is not for (tiny) children – that $(a + b) \cdot (a - b) = a \cdot a - b \cdot b$ is true for all $a, b \in \mathbb{R}$

We know – without algebra – how to test

“ $(a + b) \cdot (a - b) = a \cdot a - b \cdot b$ ”

for specific values of a and b ...

If $a = 5$ and $b = 2$, then:

$$\underbrace{\underbrace{\underbrace{a}_{5} + \underbrace{b}_{2}}_7 \cdot \underbrace{\underbrace{a}_{5} - \underbrace{b}_{2}}_3}_{21} = \underbrace{\underbrace{a}_{5} \cdot \underbrace{a}_{5}}_{25} - \underbrace{\underbrace{b}_{2} \cdot \underbrace{b}_{2}}_4}_{21}$$

true

If $a = 10$ and $b = 1$, then:

$$\underbrace{\underbrace{\underbrace{a}_{10} + \underbrace{b}_{1}}_{11} \cdot \underbrace{\underbrace{a}_{10} - \underbrace{b}_{1}}_9}_{99} = \underbrace{\underbrace{a}_{10} \cdot \underbrace{a}_{10}}_{100} - \underbrace{\underbrace{b}_{1} \cdot \underbrace{b}_{1}}_1}_{99}$$


true

A named function: $g(a) = a \cdot a + 4$

An unnamed function: $\lambda a. a \cdot a + 4$

Let $h = \lambda a. a \cdot a + 4$.

Then:



The usual notation for defining functions is like this:

$$f : \mathbb{N} \rightarrow \mathbb{R}$$

$$n \mapsto 2 + \sqrt{n}$$

$$\begin{array}{l} \text{(name)} : \text{(domain)} \rightarrow \text{(codomain)} \\ \text{(variable)} \mapsto \text{(expression)} \end{array}$$

It creates *named* functions
(with domains and codomains).

The usual notation for creating named functions
without specifying their domains and codomains
is just $f(n) = 2 + \sqrt{n}$.

Note that this is:

$$f \quad (n) \quad = \quad 2 + \sqrt{n}$$

$$\text{(name)} \quad \text{((variable))} \quad = \quad \text{(expression)}$$

The *graph* of


$$h : \begin{array}{l} \{-2, -1, 0, 1, 2\} \rightarrow \{0, 1, 2, 3, 4\} \\ k \mapsto k^2 \end{array}$$

is $\{(-2, 4), (-1, 1), (0, 0), (1, 1), (2, 4)\}$.

We can think that a function *is* its graph,
and that a lambda-expression (with domain) reduces to a graph.
Then $h = \{(-2, 4), (-1, 1), (0, 0), (1, 1), (2, 4)\}$
and $h(-2) = \{(-2, 4), (-1, 1), (0, 0), (1, 1), (2, 4)\}(-2) = 4$.

Let $h := (\lambda k : \{-2, -1, 0, 1, 2\}.k^2)$.

We have:



Note:

the graph of $(\lambda n : \mathbb{N}.n^2)$ has infinite points,
the graph of $(\lambda n : \mathbb{N}.n^2)$ is an infinite set,
the graph of $(\lambda n : \mathbb{N}.n^2)$ can't be written down *explicitly* without '...'s...

Mathematicians love infinite sets.

Computers hate infinite sets.

For mathematicians a function *is* its graph.

For computer scientists a function *is* a finite program.

Computer scientists love 'λ's!

I love things like this: $\left\{ \begin{array}{l} (3, 30), \\ (4, 40) \end{array} \right\} (3) = 30$

Types (introduction)

Let:

$$A = \{1, 2\}$$

$$B = \{30, 40\}.$$

If $f : A \rightarrow B$, then f is one of these

four functions:

$$\begin{array}{cccc} 1 \mapsto 30 & 1 \mapsto 30 & 1 \mapsto 40 & 1 \mapsto 40 \\ 2 \mapsto 30 & 2 \mapsto 40 & 2 \mapsto 30 & 2 \mapsto 40 \end{array}$$

or, in other notation,

$$\left\{ \begin{array}{l} (1,30) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,30) \\ (2,40) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,40) \end{array} \right\}$$

which means that:

$$f \in \left\{ \left\{ \begin{array}{l} (1,30) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,30) \\ (2,40) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,40) \end{array} \right\} \right\}$$

Let's use the notation " $A \rightarrow B$ " for
"the set of all functions from A to B ".

$$\text{Then } (A \rightarrow B) = \left\{ \left\{ \begin{array}{l} (1,30) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,30) \\ (2,40) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,30) \end{array} \right\}, \left\{ \begin{array}{l} (1,40) \\ (2,40) \end{array} \right\} \right\}$$

and $f : A \rightarrow B$

means $f \in (A \rightarrow B)$.

In Type Theory and λ -calculus " $a : A$ "

is pronounced " a is of type A ", and the meaning
of this is *roughly* " $a \in B$ ".

(We'll see the differences between ' \in ' and ' $:$ ' (much) later).

Note that:

1. if $f : A \rightarrow B$ and $a : A$ then $f(a) : B$
2. if $a : A$ and $b : B$ then $(a, b) : A \times B$
3. if $p : A \times B$ then $\pi p : A$ and $\pi' p : B$, where
' π ' means 'first projection' and
' π' ' means 'second projection';
if $p = (2, 30)$ then $\pi p = 2$, $\pi' p = 30$.

If $p : A \times B$ and $g : B \rightarrow C$, then:

$$\underbrace{\left(\underbrace{\underbrace{\pi p}_{:A \times B}}_{:A}, \underbrace{\underbrace{g(\pi' p)}_{:A \times B}}_{:B} \right)}_{:C} : A \times C$$

Typed λ -calculus: trees

$$A = \{1, 2\}$$

$$B = \{3, 4\}$$

$$C = \{30, 40\}$$

$$D = \{10, 20\}$$

$$A \times B = \left\{ \begin{array}{l} (1, 3), (1, 4), \\ (2, 3), (2, 4) \end{array} \right\}$$

$$B \rightarrow C = \left\{ \left\{ \begin{array}{l} (3,30), \\ (4,30) \end{array} \right\}, \left\{ \begin{array}{l} (3,30), \\ (4,40) \end{array} \right\}, \left\{ \begin{array}{l} (3,40), \\ (4,30) \end{array} \right\}, \left\{ \begin{array}{l} (3,40), \\ (4,40) \end{array} \right\} \right\}$$

If we know [the values of] a, b, f
then we know [the value of] $(a, f(b))$.

If $(a, b) = (2, 3)$ and $f = \left\{ \begin{array}{l} (3,30), \\ (4,40) \end{array} \right\}$

then $(a, f(b)) = (2, 30)$.

If we know the *types* of a, b, f
we know the type of $(a, f(b))$.

If we know the types of p, f
we know the type of $(\pi p, f(\pi' p))$.

If we know the types p, f
we know the type of $(\lambda p : A \times B.(\pi p, f(\pi' p)))$.

$$\frac{\frac{(a, b)}{a} \pi \quad \frac{\frac{(a, b)}{b} \pi' \quad f}{f(b)} \text{ app}}{(a, f(b)) \text{ pair}} \quad \frac{\frac{(2, 3)}{2} \pi \quad \frac{\frac{(2, 3)}{3} \pi' \quad \{(3, 30), (4, 40)\}}{30} \text{ app}}{(2, 30) \text{ pair}} \text{ app}$$

$$\frac{\frac{(a, b) : A \times B}{a : A} \pi \quad \frac{\frac{(a, b) : A \times B}{b : B} \pi' \quad f : B \rightarrow C}{f(b) : C} \text{ app}}{(a, f(b)) : A \times C} \text{ pair}$$

$$\frac{\frac{p : A \times B}{\pi p : A} \pi \quad \frac{\frac{p : A \times B}{\pi' p : B} \pi' \quad f : B \rightarrow C}{f(\pi' p) : C} \text{ app}}{(\pi p, f(\pi' p)) : A \times C} \text{ pair}$$

$$\frac{\frac{\frac{p : A \times B}{\pi p : A} \pi \quad \frac{\frac{p : A \times B}{\pi' p : B} \pi' \quad f : B \rightarrow C}{f(\pi' p) : C} \text{ app}}{(\pi p, f(\pi' p)) : A \times C} \text{ pair}}{(\lambda p : A \times B.(\pi p, f(\pi' p))) : A \times B \rightarrow A \times C}$$

Exercises

Let:

$$A = \{1, 2\}$$

$$B = \{3, 4\}$$

$$C = \{30, 40\}$$

$$D = \{10, 20\}$$

$$f = \left\{ \begin{array}{l} (3,30), \\ (4,40) \end{array} \right\}, f : B \rightarrow C$$

$$g = \left\{ \begin{array}{l} (1,10), \\ (2,20) \end{array} \right\}, g : A \rightarrow D$$

a) Check that:

$$(\lambda p : A \times B. (\pi p, f(\pi' p))) = \left\{ \begin{array}{l} ((1,3), (1,30)), \\ ((1,4), (1,40)), \\ ((2,3), (2,30)), \\ ((2,4), (2,40)) \end{array} \right\}$$

b) Let $p = (2, 3)$. Evaluate $(g(\pi p), f(\pi' p))$.c) Check that $(g(\pi p), f(\pi' p)) : D \times C$.d) Suppose that $p : A \times B$. Check that

$$(\lambda p : A \times B. (g(\pi p), f(\pi' p))) : A \times B \rightarrow D \times C.$$

e) Evaluate $(\lambda p : A \times B. (g(\pi p), f(\pi' p)))$.

Here is another notation for checking types:

$$\underbrace{\underbrace{\underbrace{\underbrace{\underbrace{\lambda p : A \times B. (\pi p, f(\pi' p))}_{:A \times B \rightarrow A \times C}}_{:A \times B}}_{:A} \underbrace{\underbrace{\underbrace{f(\pi' p)}_{:B \rightarrow C}}_{:B}}_{:C}}_{:A \times B}}$$

Compare it with:

$$\frac{\frac{p : A \times B}{\pi p : A} \pi \quad \frac{\frac{p : A \times B}{\pi' p : B} \pi' \quad f : B \rightarrow C}{f(\pi' p) : C} \text{pair}}{(\pi p, f(\pi' p)) : A \times C} \text{app}}{(\lambda p : A \times B. (\pi p, f(\pi' p))) : A \times B \rightarrow A \times C}$$

Represente graficamente:

$$A := \{(1, 4), (2, 4), (1, 3)\}$$

$$B := \{(1, 3), (1, 4), (2, 4)\}$$

$$C := \{(1, 3), (1, 4), (2, 4), (2, 4)\}$$

$$D := \{(1, 3), (1, 4), (2, 3), (2, 4)\}$$

$$h := \{(0, 3), (1, 2), (2, 1), (3, 0)\}$$

$$k := \{x : \{0, 1, 2, 3\}; (x, 3 - x)\}$$

$$m := \{y : \{0, 1, 2, 3\}; (3 - y, y)\}$$

$$f := (\lambda x : \{0, 1, 2\}. x \cdot x)$$

$$g := (\lambda a : \{0, 1, 2\}. 3 \cdot a)$$

$$q := (\lambda a : \{0, 1, 2\}. (x + 1)^2 - 2x - 1)$$

Tudo isto aqui a gente pode calcular com tabelas:

$$\forall x : \{0, 1, 2, 3\}. x < 2$$

$$\forall x : \{0, 1, 2, 3\}. x^2 \leq 5$$

$$\forall x : \{0, 1, 2, 3\}. x^2 < 10$$

$$\forall x : \{0, 1, 2, 3\}. x \geq 2$$

$$\exists x : \{0, 1, 2, 3\}. x^2 \geq 5$$

$$\{x : \{0, 1, 2, 3\}; x^2\}$$

$$\{x : \{0, 1, 2, 3\}, x \geq 2; x\}$$

$$\Sigma x : \{0, 1, 2, 3\}. x^2$$

$$\Pi x : \{0, 1, 2, 3\}. x + 1$$

$$\lambda x : \{0, 1, 2, 3\}. 5 - x$$

$$\forall x : \{0, 1, 2\}. P(x) = P(0) \& P(1) \& P(2)$$

$$\exists x : \{0, 1, 2\}. P(x) = P(0) \vee P(1) \vee P(2)$$

$$\{x : \{0, 1, 2\}; f(x)\} = \{f(0), f(1), f(2)\}$$

$$\Sigma x : \{0, 1, 2\}. f(x) = f(0) + f(1) + f(2)$$

$$\Pi x : \{0, 1, 2\}. f(x) = f(0) \cdot f(1) \cdot f(2)$$

$$\lambda x : \{0, 1, 2\}. f(x) = \{(0, f(0)), (1, f(1)), (2, f(2))\}$$

Obs:

$$\Sigma x : \{0, 1, 2, 2, 3\}. x^2 \rightarrow f(0) + f(1) + f(1) + f(2) \quad (?)$$

$$\Pi x : \{0, 1, 2, 2, 3\}. x + 1 \rightarrow f(0) \cdot f(1) \cdot f(1) \cdot f(2) \quad (?)$$

If $P(x) = (x < 2)$ then

$$\begin{array}{ccc} \forall x : \{0, 1, 2\}. P(x) & \longrightarrow & P(0) \& P(1) \& P(2) \\ \downarrow & & \downarrow \\ \forall x : \{0, 1, 2\}. x < 2 & \longrightarrow & (0 < 2) \& (1 < 2) \& (2 < 2) \\ & & \downarrow \\ & & 1 \& 1 \& 0 \end{array}$$